

THE TEMPORARY SOCIAL PARASITISM OF  
LASIUS SUBUMBRATUS VIERECK

---

By WILLIAM MORTON WHEELER

---

Reprinted from *Psyche*, Vol. XXIV, No. 6

# THE TEMPORARY SOCIAL PARASITISM OF *LASIUS* *SUBUMBRATUS* VIERECK.<sup>1</sup>

BY WILLIAM MORTON WHEELER.

During the past summer, while studying the ants of the Sacramento Mountains, at Cloudcroft, N. M. (alt. 9,000 ft.), I had an opportunity to observe the method of colony formation employed by *Lasius* (*Formicina*) *umbratus* Nyl. subsp. *subumbratus* Viereck. *L. umbratus* has long been known as a circumpolar species comprising several races, or subspecies and varieties, the names and known geographical distribution of which are recorded in the following list:

## PALEARCTIC FORMS.

<i>L. umbratus</i> subsp. <i>umbratus</i> Nyl.	} Northern and Alpine Eu- rasia from Britain to Japan. Russia.
subsp. <i>mixtus</i> Nyl.	
var. <i>mixto-umbratus</i> Forel	
var. <i>umbrato-affinis</i> Ruzsky	

## NEARCTIC FORMS.

subsp. <i>mixtus</i> var. <i>aphidicola</i> Walsh	Canada, Central and At- lantic States.
subsp. <i>subumbratus</i> Viereck	British America, Rocky Mts.
subsp. <i>speculiventris</i> Emery	Known only from New Jersey.
subsp. <i>vestitus</i> Wheeler	Known only from Idaho.
subsp. <i>minutus</i> Emery	Central and Atlantic States.

In Europe, according to Wasmann and Donisthorpe, the var. *mixto-umbratus* is more common than the typical *umbratus* or *mixtus*, between which it is intermediate in size, color and pilosity. In the Eastern United States *aphidicola* is certainly the most

<sup>1</sup>Contributions from the Entomological Laboratory of the Bussey Institution, Harvard University. No. 136.

abundant form of *umbratus*. The subspecies *speculiventris* and *vestitus* are imperfectly known, the latter from a single female specimen, the former only from the types taken many years ago by Pergande at Caldwell, N. J. The subspecies *minutus* is very sporadic. During the past eight years I have found only a single colony of it in the vicinity of Boston. The types of the subspecies *subumbratus* were taken by Professor Cockerell at Beulah, N. M. (alt. 8,000 ft.). I have taken it at Williams, and on Mt. Lemmon in the Santa Catalina Range (9,000 ft.), Ariz., at Cloudcroft, N. M. (9,000 ft.), Lake Tahoe, Calif. (8,000 ft.), near Ottawa, Ontario, and Hull, Quebec and have seen specimens from Digby and Bedford, Nova Scotia. It is therefore the most boreal and stenothermal of all the North American subspecies of *umbratus* and, as I have stated in a former paper (1917), properly belongs to the Canadian zone.

To the ethologist *umbratus* is of unusual interest, because in Europe it is a temporary parasite of *L. niger* and in turn serves as the host of *L. (Dendrolasius) fuliginosus*, which is therefore a true temporary social hyperparasite. The young *umbratus* queen is adopted by a colony of *niger* workers, which bring up her brood. Eventually these workers die off, probably in the course of four or five years, and the *umbratus* colony, now pure, becomes very populous. Then it may in turn adopt a queen of *fuliginosus*, the offspring of which eventually supplant the *umbratus* colony. In both cases the host queens are eliminated, probably by being assassinated either by their own workers or by the parasitic queens. The various phases in the development of the *umbratus* and *fuliginosus* colonies have been studied by European myrmecologists. The occasional occurrence of mixed colonies of *umbratus* and *niger* and *umbratus* and *fuliginosus* was noted by Schenck, Adlerz, Forel, Wasmann and de Lannoy, but Emery and Forel in 1908 first suggested that they might be interpreted as arising from temporary social parasitism. Later Crawley and Donisthorpe fully demonstrated the truth of this conjecture by showing that the typical *umbratus* is parasitic on the typical *niger*, the subspecies *mixtus* on the subspecies *alienus*, and that *umbratus* regularly functions as the temporary host of *fuliginosus*.

Although North America is so rich in *umbratus* forms, no one has hitherto been able to prove from observations in the field that any

of them is parasitic on some one of our varieties of *L. niger*. I have repeatedly found deâlated queens of *aphidicola* in small cavities under stones as if in the act of founding colonies independently in the same manner as most ants, but never with brood. Nor have I ever been able to find a mixed colony of *L. niger* var. *americanus* and *aphidicola*, even in Illinois where both forms are common in the same localities. The only indication that any American form of *umbratus* may be a social parasite is furnished by some specimens of *subumbratus* collected by Mr. W. Reiff in Nova Scotia and some experiments by Tanquary on *minutus*. At Bedford, near Halifax, Mr. Reiff took six deâlated *subumbratus* queens from three colonies of what I recorded at the time (1910) as "the large yellowish form of *Lasius niger* var. *neoniger* Emery." Since the discovery of Pergande's types of Alaskan ants I can now state definitely that this variety is *sitkaësis*. As Reiff's specimens were not accompanied by notes, the parasitism of *subumbratus* could only be conjectured. Tanquary introduced successively some eighty-eight of the small, active queens of *minutus* into twenty different colonies of *L. americanus*, *nearcticus*, *brevicornis*, *claviger* and *interjectus* and obtained one case of adoption. This was in a colony of eight workers and a large number of cocoons of *americanus*. He concludes that "although one adoption out of 88 attempts is a small percentage, yet I think the ease with which this queen was adopted is very suggestive, and taken together with the facts mentioned above, namely the sporadic occurrence of the species, the very large number of females produced, the small size of the females, the fact that these females have not been seen in the act of founding a colony and one additional fact that may be mentioned, the mimetic coloration of the females (the color of these females is exactly the same as that of the darker form of *americanus*), I think justifies us in concluding that the queen of this species is in all probability, temporarily, parasitic upon the common *L. americanus*."

My observations at Cloudcroft leave no doubt concerning the parasitism of the subspecies *subumbratus* on two forms of *L. niger*, the var. *sitkaësis* Pergande and the var. *neoniger* Emery. The former, as I have recently shown (1917), is the common form of *niger* at higher altitudes and latitudes from Alaska to Maine and Nova Scotia and southward along the ranges of the Sierra Nevada

and Rocky Mountains. The worker and female of this variety are large, brown or yellowish brown, with numerous erect or suberect hairs on the legs and antennal scapes. It passes over into the var. *neoniger*, which is smaller and darker, but has hairy legs and scapes, unlike our common and more xerothermic var. *americanus*, which closely resembles the European *alienus* in having the legs and scapes merely finely pubescent. At Cloudcroft *sitkaënsis* is everywhere abundant in the pine forests at an altitude of 9,000 to 9,500 feet, and nests by preference under large stones. The females and workers have a peculiar and rather agreeable aroma, which is much stronger than that of *americanus* and much like the odor of *subumbratus*. On rocky slopes, exposed to the sun, especially along the edges of the alpine meadows, the typical *sitkaënsis* is replaced by *neoniger* and forms intermediate in size and coloration. At altitudes of 7,000 to 7,500 feet or at higher elevations in very warm, dry situations the var. *americanus* and forms transitional to *neoniger* are not uncommon. At Cloudcroft I failed to find *subumbratus* below 9,000 feet. Its colonies are very populous and live under large stones, usually in the shade of the pines. Its habits, like those of the other subspecies of *umbratus*, are decidedly hypogæic. It does not make carton like the European *umbratus*. This is generally true also of our eastern *aphidicola*. I have, however, found a few colonies of this ant at Colebrook, Conn., and Ottawa, Ontario, nesting in dark brown carton nests in the centers of rotten logs.

The observations proving that the young queens of *subumbratus* establish their colonies with the aid of *sitkaënsis* and *neoniger* workers may now be transcribed from my note-book:

*July 3 and 5.* Four colonies of *subumbratus* were found under large stones. Two were of considerable size and the superficial chambers of the nests were full of black males and golden yellow females ready for the marriage flight.

*July 8.* The marriage flight of *subumbratus* must have occurred on July 6 or 7 as there were numerous deälated females running about among the stones on an open, sunny slope where many of the stones covered nests of *neoniger*. My son Ralph assisted me in carefully turning over the stones and examining these nests. In one area, about 200 feet in diameter, nearly every nest had from one to five deälated *subumbratus* queens running about in the large

shallow superficial chambers. By shading the chambers, which were occupied by numerous *neoniger* workers and cocoons, the behavior of the yellow queens could be readily studied. They ran in and out of the chambers and galleries leading into the soil and, when accosted by the *neoniger* workers, made peculiar supplicatory movements with the antennæ, precisely like those of *Formica consocians* queens seeking adoption in nests of *F. incerta*. The *neoniger* workers, however, resented the intrusion and frequently seized the legs and antennæ of the queens, though they soon released their hold. The queens thus liberated often escaped from the nest and hid under stones near it or even in unfrequented nooks in the superficial chambers of the nest itself. On several occasions we saw a queen seize a *neoniger* worker, carry it out of the nest and release it uninjured. One queen was seen to snatch up a worker cocoon, carry it out of the nest and conceal it under an adjacent pebble. This is very suggestive of the behavior of the young *Formica sanguinea* queen that has entered a nest of *F. fusca*, but though it undeniably shows an interest on the part of the *subumbratus* queen in the *neoniger* brood, her method of colony formation is not that of *F. sanguinea*, for we soon discovered a depauperate *neoniger* colony which had undoubtedly adopted a *subumbratus* queen. The latter was moving to and fro very slowly and amicably in the midst of several *neoniger* workers and their cocoons in a small recess under a stone. The ants were watched for some time and behaved precisely like members of a single colony. That the queens are nevertheless occasionally treated much more severely by the *neoniger* workers, is apparent from the fact that we found a few recently killed and one partially eaten in the superficial chambers of some of the nests.

July 9. Visited another rocky slope and an adjoining pine forest about a mile east of Cloudercroft and two miles from the locality of yesterday's observations. Under the stones covering *neoniger* and *sitkaënsis* nests there were many dealated *subumbratus* queens eagerly seeking adoption. The following three colonies showed clearly that some of the queens had been or were being accepted:

1. A populous *sitkaënsis* colony under several large contiguous stones among the pines. The superficial galleries of the nest were filled with workers and worker, male and female cocoons nearly

ready to hatch. Under two of the stones a couple of *subumbratus* queens was found, each lurking quietly in a little cavity at the edge of a mass of cocoons nearly filling a large, shallow superficial chamber.

2. A similar but even larger *sitkaënsis* nest, also under a pile of stones, under three of which *subumbratus* queens were found. One of the queens was in a small cavity at the edge of a heap of cocoons while each of the others was resting on the very center of a collection of cocoons nearly as large as my palm. These queens had certainly been adopted and the picture presented was precisely like that of the recently adopted *Formica consocians* queens mounted on the cocoons and surrounded and fondled by the workers of *F. incerta* (Wheeler 1906). When the nest was disturbed the queens hurried into the galleries and disappeared in the soil. They had evidently become strongly negatively heliotactic, unlike queens that are still running about on the soil immediately after the marriage flight.

3. A fine *sitkaënsis* nest under a single large stone among the pines had an adopted *subumbratus* queen surrounded by dozens of workers in the center of a pile of worker and female cocoons nearly as large as my hand. She quickly slipped into a burrow and was only recovered by digging down into the soil to a depth of more than six inches. In this nest a few of the female *sitkaënsis* pupæ were naked, *i. e.*, not enclosed in cocoons.

Further search in the same locality brought to light two mixed colonies consisting of *subumbratus* and *sitkaënsis* workers. I failed to find the queens of the former species, which were probably present but hiding deep in the soil. The failure to find them is not surprising as one very rarely secures the mother queen of a fully developed colony of any of our species of *Lasius* without extensive and very careful excavation of the nest.

The foregoing observations show that the *subumbratus* queen has to acquire the brood odor of the host before she can be accepted. She therefore lurks very near the cocoons in a small earthen cavity and somewhat later, in the early stage of adoption, like *Formica consocians*, takes up a position on the brood. This renders her immune to attack by the host workers and gradually accustoms them to her presence. That adoption can be secured in very populous and flourishing colonies of *sitkaënsis* is shown by the

observations of July 9. Nevertheless these observations throw no light on three interesting problems, which may be briefly considered:

1. The first problem relates to the elimination of the host queen, evidently a *conditio sine qua non* of successful colony formation by such temporary social parasites as *subumbratus*. The elimination is supposed to be effected in one of three ways: Either adoption succeeds only in host colonies that have already lost their rightful queen through death by accident or old age, or the queen is killed by her own workers after and because the young parasitic queen has secured their allegiance, or the latter assassinates the host queen. The first hypothesis seems to be supported by the above described adoption of a *subumbratus* queen in a depauperate colony of *neoniger*, although it is by no means certain that this colony was queenless. The large *sitkaënsis* colonies containing recently adopted *subumbratus* queens, however, would seem to support the second or third hypothesis. In regard to this problem in the European *umbratus*, Donisthorpe (1915) says: "No cases are known of the host queen and the parasite living together in a nest, so, unless a female can only be adopted by a queenless colony, it must sometimes happen that a female is accepted by a colony already possessing a queen of its own species. In such a case the intruder must either kill the rightful queen herself, as the female of (non-British) *Bothriomyrmex* kills the queen of her host *Tapinoma* or the workers of the host species must themselves assassinate their own queen, as do the workers of *Tetramorium caespitum* when they have accepted a female *Anergates atratulus*." He adds, however, that "Crawley confined several queens of *umbrata* with queens of *nigra* and the latter were always killed by the former, which although a little the smaller, is stronger and possesses more powerful mandibles." It seems to me that the morphological peculiarities suggested by Donisthorpe, namely the much broader and larger head and stronger mandibles so characteristic of the female of all *umbratus* forms as compared with the same sex of *niger*, indicate very clearly that the parasite actually decapitates the host queen in much the same manner as *Bothriomyrmex decapitans* was seen by Santschi (1906) to decapitate the *Tapinoma erraticum* queen and the queen *Wheeleriella santschii*, the queen of *Monomorium solomonis* (Forel 1906).



2. The fact that one of the *sitkaënsis* colonies observed July 9 contained two fully adopted *subumbratus* queens and one on the verge of adoption suggests the question as to whether all three queens would survive as mothers of the compound colony or whether two of them would be subsequently eliminated. This question can, of course, be answered only by further investigation. But as the various species of the genus *Lasius* hitherto observed are decidedly haplometrotic it seems probable that at least in the majority of cases of adoption of several *subumbratus* queens by a *sitkaënsis* or *neoniger* colony, the number must be subsequently reduced to unity either by combats among the queens or by the regulatory intervention of the workers.

3. My field observations leave some doubt as to whether *sitkaënsis* or *neoniger* is the true, or normal host of *subumbratus*. Indirect evidence, however, points to the former variety, since the queens and workers of *subumbratus* are much more like the corresponding phases of *sitkaënsis* in size and coloration and since pure adult colonies of these forms prefer the same shady environment and have the same hypogæic habits. Reiff's Nova Scotia specimens of *subumbratus* queens from *sitkaënsis* nests may also be cited in this connection. *L. neoniger*, therefore, would be an unusual or perhaps recently acquired host, and the conditions would be comparable to those of the typical European *umbratus*, whose normal host is the typical *niger*, though it is occasionally parasitic, as Donisthorpe has shown (1915), on the subspecies *alienus*, the normal host of *mixtus*.

The rather large number of observed cases of adoption of *subumbratus* seems to indicate that the queen of this subspecies is specially favored as compared with the queens of some other temporary social parasites (e. g. *Formica rufa* and *truncicola*) by the possession of peculiar myrmecophilous characters (trichomes, exudate organs). I believe that the beautiful golden yellow color and pilosity of the *subumbratus* queen, which make it resemble the queens of *Formica consocians*, *ciliata*, *criniventris* and *oreas*, the smaller stature and the delicate odor may be designated as such characters. In this connection I may call attention to the queen of *Lasius crinitus*, long ago described by Frederick Smith from Cashmir, as being in all probability a temporary social parasite on *L. niger* or one of its varieties. I have recently seen a few queens

of *crinitus* from Sikkim and was struck by their remarkable resemblance in color and pilosity to the queens of the Colorado *F. ciliata*. The source of the delicate odor of *subumbratus*, of the strong, lemon-verbena odor of our North American *Lasii* of the subgenus *Acanthomyops* and of the peculiar pungent, sweetish odor of the Eurasian *fuliginosus*, is worthy of further investigation. Meinert and Bönner believe that in *umbratus* and *fuliginosus* it comes from the salivary glands. Forel had previously traced it to the head as this portion of the body when crushed emitted a more violent aroma than the thorax and gaster. I have been under the impression that it might originate in the epinotal glands. Since it is so powerful and pervasive that it even saturates the walls of the nest a renewed and careful investigation may be required before its precise source in the body of the insect is known.

#### LITERATURE.

1886. Adlerz, G. Myrmekologiska Studier II. Svenska Myror och deras Lefnadsförhållanden. Bih. K. Svensk. Vet.-Akad. Handl. 11, 18, 1886, pp. 1-329, 7 pls.
1915. Bönner, W. Der temporäre soziale Hyperparasitismus von *Lasius fuliginosus* und seine Beziehungen zu *Claviger longicornis* Müll. Zeitschr. f. wiss. Insekt. Biol. 11, 1915, pp. 14-20.
1909. Crawley, W. C. Queens of *Lasius umbratus* Nyl., Accepted by Colonies of *Lasius niger* L. Ent. Month. Mag. (2) 20, 1909, pp. 94-99.
1910. Crawley, W. C. Workers of *Lasius flavus* (? *L. umbratus*) among *Lasius fuliginosus*. Ent. Rec. 22, 1910, pp. 67-69.
1913. Crawley, W. C. and Donisthorpe, H. St. J. K. The Founding of Colonies of Queen Ants. Trans. Second Intern. Ent. Congr. (1912) 1913, 2, pp. 11-77.
1911. Donisthorpe, H. St. J. K. Further Observations on Temporary Social Parasitism and Slavery in Ants. Trans. Ent. Soc. London, 1911, pp. 175-183.
1913. Donisthorpe, H. St. J. K. On Some Remarkable Associations between Ants of Different Species. Rep. Lancast. Ches. Ent. Soc. 36 (1912), 1913, pp. 38-56.
1915. Donisthorpe, H. St. J. K. British Ants, Their Life-History

- and Classification. Plymouth, William Brendon & Son. 1915.
1911. Donisthorpe, H. St. J. K. and Crawley, W. C. Experiments on the Formation of Colonies of *Lasius fuliginosus* Females. Trans. Ent. Soc. London, 1912, pp. 664-672.
1908. Emery, C. Remarques sur l'existence de *Lasius mixtus* dans les fourmilières de *L. fuliginosus*. Ann. Soc. Ent. Belg. 52, 1908, pp. 182-183.
1906. Forel, A. Mœurs des Fourmis parasites des Genres *Wheeleria* et *Bothriomyrmex*. Rev. Suisse Zool. 14, 1906, pp. 51-69, 6 text-figs.
1908. Forel, A. Lettre à la Société Entomologique de Belgique. Ann. Soc. Ent. Belg. 52, 1908, pp. 180-181.
1908. de Lannoy, F. Notes sur le *Lasius niger* et le *Lasius fuliginosus*. Ann. Soc. Ent. Belg. 52, 1908, pp. 47-53.
1906. Santschi, F. À propos des mœurs parasitiques temporaires des Fourmis du genre *Bothriomyrmex*. Ann. Soc. Ent. France, 75, 1906, pp. 363-392.
1852. Schenck, A. Beschreibung nassauischer Ameisenarten. Jahrb. Ver. Naturk. Nassau, 8, 1, 1852, pp. 3-149.
1911. Tanquary, M. Experiments on the Adoption of *Lasius*, *Formica* and *Polyergus* Queens by Colonies of Alien Species. Biol. Bull. 20, 1911, pp. 281-308.
1908. Wasmann, E. Weitere Beiträge zum sozialen Parasitismus und der Sklaverei bei den Ameisen. Biol. Central. 28, 1908, pp. 257-271, 289-306, 321-333, 353-382, 417-441.
1909. Wasmann, E. Ueber gemischte Kolonien von *Lasius*-Arten. Zool. Anzeig. 35, 1909, pp. 129-141.
1910. Wasmann, E. Nachträge zum sozialen Parasitismus und der Sklaverei bei den Ameisen. Biol. Centralbl. 30, 1910, pp. 453-464, 475-496, 515-524.
1906. Wheeler, W. M. On the Founding of Colonies by Queen Ants, with Special Reference to the Parasitic and Slave-making Species. Bull. Amer. Mus. Nat. Hist. 22, 1906, pp. 33-105, 7 pls.
1910. Wheeler, W. M. The North American Forms of *Lasius umbratus* Nylander. Psyche, 17, 1910, pp. 235-243.
1917. Wheeler, W. M. The Mountain Ants of Western North America. Proc. Amer. Acad. Sci. 52, 1917, pp. 457-569.